

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

12

EUROPEAN PATENT APPLICATION

21 Application number: 84304888.5

22 Date of filing: 18.07.84

51 Int. Cl.: **C 07 D 401/12, C 07 D 487/04,**
C 07 D 473/10, C 07 D 471/04,
A 61 K 31/44
// C07D211/90

30 Priority: 23.07.83 GB 8319886
07.01.84 GB 8400354

43 Date of publication of application: 30.01.85
Bulletin 85/5

84 Designated Contracting States: AT BE CH DE FR GB IT
LI LU NL SE

71 Applicant: **Pfizer Limited, Ramsgate Road, Sandwich
Kent CT13 9NJ (GB)**

84 Designated Contracting States: GB

71 Applicant: **Pfizer Corporation, Calle 15 1/2 Avenida
Santa Isabel, Colon (PA)**


84 Designated Contracting States: BE CH DE FR IT LI LU
NL SE AT

72 Inventor: **Alker, David, Dr., 21 Whitewood Road, Eastry,
Nr. Deal Kent (GB)**
Inventor: **Cross, Peter Edward, Dr., 21 Cherry Avenue,
Canterbury Kent (GB)**
Inventor: **Campbell, Simon Fraser, Dr., Grey Friars Upper
Street, Kingsdown, Deal Kent (GB)**

74 Representative: **Moore, James William, Pfizer Limited
Ramsgate Road, Sandwich Kent CT13 9NJ (GB)**

54 Dihydropyridine anti-Ischaemic and antihypertensive agents.

57 1,4-Dihydropyridine derivatives of the formula:



(I)

wherein R is aryl or heteroaryl; R¹ and R² are each independently C₁-C₄ alkyl or 2-methoxyethyl; X is a 5 or 6 membered nitrogen-containing heterocyclic ring which is substituted with one or more hydroxyl or oxo groups and which may optionally be fused to a further 5 or 6 membered nitrogen-containing heterocyclic ring, and which may optionally be further substituted in the heterocyclic ring or further fused heterocyclic ring; Y is -(CH₂)_n-, -CH₂CH(CH₃)- or -CH₂C(CH₃)₂-; and n is 1 to 3 when X is linked to Y by a ring carbon atom, or 2 or 3 when X is linked to Y by a ring nitrogen atom; and their pharmaceutically acceptable salts, and pharmaceutical preparation containing such compounds, have utility as anti-ischaemic and antihypertensive agents.


This invention relates to certain dihydropyridines,
specifically to certain 1,4-dihydropyridines having a hydroxy or
oxo-substituted heterocyclic ring in a side chain attached to the
2-position, which have utility as anti-ischaemic and
5 antihypertensive agents, and to pharmaceutical preparations
containing such compounds.

The compounds of the invention reduce the movement of calcium
into the cell and they are thus able to delay or prevent the
cardiac contracture which is believed to be caused by an
10 accumulation of intracellular calcium under ischaemic conditions.
Excessive calcium influx during ischaemia can have a number of
additional adverse effects which would further compromise the
ischaemic myocardium. These include less efficient use of oxygen
for ATP production, activation of mitochondrial fatty acid
15 oxidation and possibly, promotion of cell necrosis. Thus the
compounds are useful in the treatment or prevention of a variety
of cardiac conditions, such as angina pectoris, cardiac
arrhythmias, heart attacks and cardiac hypertrophy. The compounds
also have vasodilator activity since they can inhibit calcium
20 influx in cells of vascular tissue and they are thus also useful
as antihypertensive agents and for the treatment of coronary
vasospasm.

According to the specification of our European Patent
Application no. 60674 there are described and claimed a number of
25 dihydropyridine anti-ischaemic and antihypertensive agents wherein
the 2-position of the dihydropyridine ring is substituted with
certain N,N-disubstituted-aminoalkoxymethyl groups. Our

co-pending European patent application no. 100189 describes and claims a related series of compounds wherein the 2-position is substituted with an aromatic heterocyclalalkoxymethyl group. We have now discovered a further series of dihydropyridine compounds
 5 having valuable therapeutic properties wherein the 2-position substituent bears a hydroxy- or oxo-substituted heterocyclic group.

Thus according to the invention, there are provided novel 1,4-dihydropyridine derivatives of the formula:-



and their pharmaceutically acceptable salts;

wherein R is aryl or heteroaryl;

R¹ and R² are each independently C₁-C₄ alkyl or 2-methoxyethyl;

15 X is a 5 or 6 membered nitrogen-containing heterocyclic ring which is substituted with one or more hydroxyl or oxo groups and which may optionally be fused to a further 5 or 6 membered nitrogen-containing heterocyclic ring, and which may optionally be additionally
 20 substituted in the heterocyclic ring or further fused heterocyclic ring by one or more C₁-C₄ alkyl, phenyl, CN, N(R³)₂, (CH₂)_mCO₂R³ or (CH₂)_mCON(R³)₂ groups,

wherein each R^3 is independently H or C_1-C_4 alkyl or the two groups R^3 are taken together with the nitrogen atom to which they are attached to form a pyrrolidinyl, piperidino, morpholino, piperazinyl or N-(C_1-C_4 alkyl)-piperazinyl group, and m is 0 or 1;

5 Y is $-(CH_2)_n-$, $-CH_2CH(CH_3)-$ or $-CH_2C(CH_3)_2-$;

and n is 1 to 3 when X is linked to Y by a ring carbon atom, or 2 or 3 when X is linked to Y by a ring nitrogen atom.

The pharmaceutically acceptable acid addition salts of the compounds of the formula (I) are those formed from acids which form non-toxic acid addition salts, for example the hydrochloride, hydrobromide, sulphate or bisulphate, phosphate or acid phosphate, acetate, citrate, fumarate, gluconate, lactate, maleate, succinate and tartrate salts. For compounds with an acidic substituent

10 salts may also be formed with bases, examples are the sodium, potassium and ammonium salts.

The term "aryl" as used in this specification for R, includes phenyl and phenyl substituted by one or two substituents each independently selected from nitro, halo, C_1-C_4 alkyl, C_1-C_4 alkoxy, hydroxy, trifluoromethyl and cyano. It also includes 1-

20 and 2-naphthyl.

The term "heteroaryl" as used in this specification for R means an aromatic heterocyclic group which may optionally be substituted and includes, for example, benzofuranyl; benzothienyl; pyridyl optionally substituted by methyl, methylthio, cyano or

25 halo; quinolyl; benzoxazolyl; benzothiazolyl; furyl; pyrimidinyl; thiazolyl; 2,1,3-benzoxadiazol-4-yl; 2,1,3-benzothiadiazol-4-yl; and thienyl optionally monosubstituted by halo or C_1-C_4 alkyl.

The 5 or 6 membered nitrogen-containing heterocyclic ring X may contain from one to three nitrogen atoms in the ring and may be saturated or unsaturated. As will be understood by those skilled in the art the hydroxyl- or oxo-substituted heterocyclic rings can be subject to keto-enol tautomerism and the group may be present either as a free hydroxyl group when in the tautomeric enol form, or as an oxo group when in the keto form. Which particular isomer is present in normal circumstances can be readily determined by appropriate physical measurements, such as for example by infra-red spectroscopy, and, in some instances the compounds may exist as mixtures of the two forms. However it is to be understood that the invention includes both forms of the compounds where they can exist and, in the description and Examples hereto, references to the hydroxy or keto form of the compound include all possible tautomeric forms and are not necessarily to be taken to indicate which is the predominant tautomeric form of the compound.

The ring X may be linked to the group Y either by a ring carbon or a ring nitrogen atom, with the proviso that, if X is linked by a ring nitrogen atom, Y must contain at least two carbon atoms (i.e. n is 2 or 3).

Examples of suitable heterocyclic rings include 2-, 4- and 5-imidazolone, 3-pyrazolone, pyrroline-2,5-dione, imidazolidine-2,4-dione, 2-pyridone, 3-piperazinone, 4-pyrimidone, pyrimidine-2,4-dione, 5-hydroxy-(1H)-1,2,3-triazole and 1-oxo-1,3-dihydropyrrolo[2,3b]pyridine.

The heterocyclic ring X may contain further substituents.

Preferred substituent groups are amino, lower alkyl, particularly methyl, carbamoyl and CN. Thus particular and preferred examples of X include 1-(2-amino-4-imidazolone),

- 5 1-(imidazolidine-2,4-dione), 1-(4-amino-2-imidazolone),
2-(1-oxo-1,3-dihydropyrrolo[2,3b]pyridine), 6-(2-amino-4-pyrimidone) and 6-(2,3-dimethyl-4-pyrimidone).

The term "halo" means fluoro, chloro, bromo or iodo. Alkyl and alkoxy groups having 3 or more carbon atoms can be straight or
10 branched chain. Compounds containing asymmetric centres will exist as one or more pairs of enantiomers and the invention includes the separated d- and l- optically active isomers as well as mixtures thereof.

Preferred values for R are 2-chlorophenyl and
15 2,3-dichlorophenyl. R^1 and R^2 are preferably CH_3 or C_2H_5 ; the case where R^1 is CH_3 and R^2 is C_2H_5 being especially preferred. Y is preferably $-(CH_2)_2-$ when X is linked to Y by a ring nitrogen atom, or $-CH_2-$ when X is linked to Y by a ring carbon atom. Thus particular and preferred examples of compounds of the invention
20 include the following:

2-Amino-1- $\left\{2- < [4-(2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy > ethyl \right\} -$
4-imidazolone;

1- $\left\{2- < [4-(2,3-Dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy > ethyl \right\} -$
25 imidazolidine-2,4-dione;

4-Amino-1- { 2- < [4- (2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy > ethyl } -
2-imidazolone;

2- { 2- < [4- (2,3-Dichlorophenyl)-3-ethoxycarbonyl-5-methoxy-
5 carbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy > ethyl } -1-oxo-
1,3-dihydropyrrolo[2,3-b]pyridine;


6- { [4- (2,3-Dichlorophenyl)-3-ethoxycarbonyl-5-methoxy-
carbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxymethyl }
-2,3-dimethyl-4-pyrimidone, and

10 2-Amino-6- { [4- (2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxymethyl } -4-pyrimidone

the latter compound being particularly preferred as having especially favourable properties.

15 The compounds of the invention can be prepared by a number of different processes according to the invention.

(a) In one process the compounds of formula I can be prepared by the Hantzsch synthesis, according to the following reaction scheme:



wherein R, R¹, R², X and Y are as previously defined.


In a typical procedure, the ketoester (III) and aldehyde are
 5 heated under reflux in a suitable organic solvent, e.g. a C₁-C₄
 alkanol such as ethanol, for about 15 minutes, and then the amino-
 crotonate (II) is added. Alternatively the aminocrotonate (II),
 the ketoester (III) and the aldehyde can be heated together in the
 solvent. Preferably a small amount of a lower alkanolic acid such
 10 as acetic acid is added to neutralise the solution. The resulting
 solution can then be heated at 60°-130°C, preferably under reflux,
 until the reaction is essentially complete, typically in 24 hours
 or less. The product of the formula (I) is then isolated and
 purified by conventional procedures, for example by partition,
 15 recrystallisation or by chromatography.

The ketoesters (III) are either known compounds or they can be prepared by methods analogous to those of the prior art, such as for example the method of Troostwijk and Kellogg, J.C.S. Chem. Comm., 1977, page 932, or using the methods illustrated in

- 5 European Patent application 60674. Similarly the amino-crotonates (II) and the aldehydes are either known or can be prepared by known methods in accordance with literature precedents.

(b) In an alternative variation of the Hantzsch synthesis the compounds of formula (I) can be prepared by the following

10 process:-




- The crotonate (V) is typically prepared in situ by reaction of the corresponding ketoester (III) with ammonium acetate by refluxing in a suitable organic solvent, e.g. a C₁-C₄ alkanol such as ethanol, for up to an hour. The crotonate (V) is then reacted with compound (IV), typically by heating in the solvent for up to about 5 hours at temperatures of from 60° to 130°C. The product (I) can then be isolated and purified by conventional procedures as before.
- 15


The starting materials (IV) are again either known compounds or they may be prepared by methods analogous to those of the prior art in accordance with literature precedents, see for example Can. J. Chem., 1967, 45, 1001.

5 It will be appreciated that while all the compounds of the invention may be prepared by the processes described under (a) or (b) above, in many cases particular compounds may more conveniently be prepared starting with the preformed dihydropyridine. Such processes are within the knowledge and
10 skill of those versed in the art and will vary according to the nature of the heterocyclic ring X desired. The following processes describe examples of a number of processes for preparing the compounds of formula (I) containing particular heterocyclic rings but other alternatives and variations will be readily
15 evident to those skilled in the art.

(c) Thus in a further process compounds of the formula (I) wherein X is a 4-pyrimidone or 3-pyrazolone group may be prepared from the corresponding dihydropyridine ketoester of formula:



20 wherein R, R¹, R² and Y are as previously defined and R⁴ is C₁-C₄ lower alkyl, by reacting either with a compound of the formula:



wherein R^5 is $\text{N}(\text{R}^3)_2$, $\text{C}_1\text{-C}_4$ alkyl or phenyl, where R^3 is as previously defined, to give compounds wherein X is a 2-substituted-4-pyrimidone group; or with hydrazine to give
5 compounds wherein X is a 3-pyrazolone group.

Thus reaction of the compound of formula (VI) with guanidine (VIa; $\text{R}^5 = \text{NH}_2$) yields the corresponding compound of formula (I) wherein X is 2-amino-4-pyrimidon-6-yl. Similarly reaction with acetamidine (VIa; $\text{R}^5 = \text{CH}_3$) gives the corresponding compound of
10 formula (I) wherein X is 2-methyl-4-pyrimidon-6-yl. The reaction in each case is performed with the reactants dissolved in an organic solvent, e.g. ethanol, in the presence of an organic base. Several days at room temperature may be required before the reaction is substantially complete and the product is then
15 isolated conventionally using normal extraction, washing and evaporation techniques, and purified, if necessary, by recrystallisation or by chromatography.


The reaction can also be performed in a similar manner using hydrazine hydrate to give the compound of formula (I) wherein X is
20 a 5-(3-pyrrazolone) group.

A further variant of this process can also be used to prepare certain compounds of the formula (I) wherein X is a bicyclic system incorporating a pyrimidone ring. Thus, for example, reaction of a compound of the formula (VI) with 2-aminoimidazolium sulphate gives the compound of formula (I) wherein X is a 7-(5-hydroxyimidazo[1,2-a]pyrimidine) group. Similarly, reaction with 2-iminopiperidine gives the compound where X is a 4-oxo-tetrahydropyrido[1,2-a]pyrimidine group.

Simple transformation reactions can of course be applied to the final products to introduce or modify substituent groups in accordance with conventional and well known procedures. Thus methylation, for example using iodomethane in the presence of a base, can be used to prepare the compound of formula I wherein X is 2-amino-3-methyl-4-pyrimidone from the corresponding 2-amino-4-pyrimidone. Similarly treatment of the compound of formula I wherein X is 2-methyl-4-pyrimidone with iodomethane yields the corresponding compound where X is 2,3-dimethyl-4-pyrimidone.

The ketoester starting materials of formula (VI) are prepared from the corresponding (1,4-dihydropyridin-2-yl)methoxy alkanoic acid by standard procedures. Thus one route that we have employed is to react the imidazolidine derivative of the acid with the sodium salt prepared from 2,2-dimethyl-1,3-dioxane-4,6-dione with sodium hydride. Ring opening of the product is effected by heating in a lower alkanol to yield the desired ketoester. The amidines and guanidines of formula (VIa) are known compounds.


(d) In a further process, compounds wherein X is a 1-(2-amino-4-imidazolone) group are prepared from a compound of the formula:



5 wherein R, R¹, R², R⁴ and Y are as previously defined, by reacting with ammonia. The reaction is readily performed by adding the compound of formula (VII) to a saturated solution of methanolic ammonia. After several hours at room temperature the solvent is evaporated to yield the desired product.

10 The starting materials of formula (VII) are prepared from the corresponding 2-[(N-alkoxycarbonylmethyl)amino]alkoxymethyl-1,4-dihydropyridines by reacting with cyanogen bromide in chloroform for several hours at room temperature.

(e) In a further process, compounds wherein X is a 1-
15 (imidazolidine-2,4-dione) group or a 1-(4-amino-2-imidazolone) group which may be substituted on the 3- or 5- positions with lower alkyl groups are prepared from compounds of the formula:




wherein R, R¹, R² and Y are as previously defined, each R⁶ is independently H or C₁-C₄ alkyl and Q is CO₂(C₁-C₄ alkyl) or CN; by reaction with a base.

- 5 The base catalysed ring closure occurs readily and is conveniently achieved by, for example, adding sodium hydride to a suspension of the compound (VIII) in a lower alkanol, and stirring for one to two hours at room temperature. The solvent is removed and the product purified by conventional techniques. Thus
- 10 reaction of the compound of formula (VIII) wherein each R⁶ is hydrogen and Q is CO₂C₂H₅ with base yields the compound of formula (I) wherein X is imidazolidin-2,4-dion-1-yl. Similarly reaction of the compound wherein each R⁶ is hydrogen and Q is CN gives the compound of formula (I) wherein X is 4-amino-2-imidazolone-1-yl.

- 15 The urea starting materials of formula (VIII) wherein Q is CO₂(C₁-C₄ alkyl) are prepared as described in our co-pending European Patent application no. 84301452.3 from the corresponding 2-[(N-alkoxycarbonylmethyl)amino]alkoxymethyl)-dihydropyridine by reaction with an isocyanate.

The compounds of formula (VIII) wherein Q is CN are prepared from the corresponding 2-[(N-cyanomethyl)amino]alkoxymethyl-1,4-dihydropyridines by reaction with isocyanates.

(f) The compounds of formula (I) wherein X is a 1-(5-amino-2-imidazolone) or a 1-(imidazolidine-2,5-dione) group are prepared
5 in a similar manner to the processes described under (d) and (e) above but starting with a compound of the formula:




wherein R, R¹, R², Q and Y are as previously defined.

10 Thus in the case where Q is CO₂(C₁-C₄ alkyl), the cyclisation is effected by stirring the compound of formula (IX) in methanolic ammonia at room temperature for several hours, typically overnight, to yield the product wherein X is imidazolidin-2,5-dione-1-yl.

15 Alternatively, when Q is CN, the cyclisation can be effected for example in methanol with sodium methoxide at room temperature for an hour or so, to yield the compound of formula (I) wherein X is a 1-(5-amino-2-imidazolone) group.


The starting materials of formula (IX) are again prepared from the corresponding 2-(aminoalkoxymethyl)-1,4-dihydropyridines by reaction either with carbonyldiimidazole followed by reacting with aminoacetonitrile to give the compound of formula (IX) wherein Q is CN, or with an alkoxycarbonylmethyl isocyanate to give the compound wherein Q is $\text{CO}_2(\text{C}_1\text{-C}_4 \text{ alkyl})$.

(g) The compounds of formula (I) wherein X is a 2-methyl-5-imidazolone group are prepared from the corresponding 2-(aminoalkoxymethyl)-1,4-dihydropyridine by reacting with ethyl N-(ethoxycarbonylmethyl)acetimidate according to the following reaction scheme:



The reaction occurs at room temperature and is conveniently effected by stirring a solution of the reactants in a lower alkanol solvent, e.g. ethanol, for several hours.

(h) The compounds of formula (I) wherein X is a 4-ethoxycarbonyl-5-hydroxy-(1H)-1,2,3-triazole group are prepared from the corresponding 2-azidoalkoxymethyl-1,4-dihydropyridine of formula:



5 wherein R, R¹, R² and Y are as previously defined by reacting with diethyl malonate in the presence of a base. The reaction is conveniently effected by heating the reactants in a lower alkanolic solvent to which sodium hydride has been added and a period of up to 24 hours under reflux is generally sufficient to
10 complete the reaction. The product is isolated and purified using conventional procedures.

The azido starting materials of formula (XI) and amines of formula (X) are prepared as described in European Patent Application Publication no. 89167.

15 1) The compounds of formula (I) wherein X is a 2-(1-oxo-1,3-dihydropyrrolo[2,3-b]pyridine) group are prepared from the corresponding 2-(aminoalkoxymethyl)-1,4-dihydropyridines (X) by reacting with 2-bromomethyl-3-ethoxycarbonylpyridine in an inert

organic solvent, e.g. acetonitrile in the presence of an acid scavenger e.g. potassium carbonate. A period of one or two days under reflux may be necessary to complete reaction and the product is then isolated and purified by conventional techniques.

5 j) The compounds of formula (I) wherein X is a 1-(5-cyano-3-methylpyrimidine-2,4-dione) group are prepared from the corresponding 2-(aminoalkoxymethyl)-1,4-dihydropyridines (X) by reaction with 2-cyano-N-ethoxycarbonyl-2-ethoxymethylene-N-methylacetamide in a lower alkanol solvent in the presence of a
10 trace of base. A period of 2 or 3 hours refluxing in methanol is generally sufficient.

k) The compounds of formula (I) wherein X is a 1-(dihydro-pyrrol-2,5-dione) group are prepared from the corresponding 2-(aminoalkoxymethyl)-1,4-dihydropyridines (X) by
15 reacting with maleic anhydride in an inert organic solvent e.g. acetonitrile under reflux for one or two hours.

The ability of the compounds to inhibit the movement of calcium into the cell is shown by their effectiveness in reducing the response of isolated heart tissue to an increase in calcium
20 ion concentration in vitro. The test is performed by mounting spirally cut strips of rat aorta with one end fixed and the other attached to a force transducer. The tissue is immersed in a bath of physiological saline solution containing potassium ions at a concentration of 45 millimolar and no calcium. Calcium chloride

is added to the bath with a pipette to give a final calcium ion concentration of 2 millimolar. The change in tension caused by the resulting contraction of the tissue is noted. The bath is drained and replaced with fresh saline solution and after 45
5 minutes, the procedure is repeated with the particular compound under test present in the saline solution. The concentration of compound required to reduce the response by 50% is recorded.

The antihypertensive activity of the compounds is also evaluated after oral administration by measuring the fall in blood
10 pressure in spontaneously hypertensive rats or renally hypertensive dogs.

For administration to man in the curative or prophylactic treatment of cardiac conditions and hypertension, oral dosages of the compounds will generally be in the range of from 2-100 mg
15 daily for an average adult patient (70 kg). Thus for a typical adult patient, individual tablets or capsules contain from 1 to 10 mg of active compound, in a suitable pharmaceutically acceptable vehicle or carrier. Dosages for intravenous administration would typically be within the range 1 to 10 mg per single dose as
20 required. In practice the physician will determine the actual dosage which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case but there can, of course, be individual instances where higher or
25 lower dosage ranges are merited, and such are within the scope of this invention.

For human use, the compounds of the formula (I) can be administered alone, but will generally be administered in admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. For example, they may be administered orally in the form of tablets containing such excipients as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs or suspensions containing flavouring or colouring agents. They may be injected parenterally, for example, intravenously, intramuscularly or subcutaneously. For parenteral administration, they are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood.

Thus in a further aspect the invention provides a pharmaceutical composition comprising a compound of the formula (I), or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable diluent or carrier.

The invention also provides a compound of the formula (I) or a pharmaceutically acceptable salt thereof, for use in the treatment of cardiovascular conditions including use in the prevention or treatment of cardiac conditions, or use as an antihypertensive, in man.

The preparation of the compounds of the invention is illustrated by the following Examples.

EXAMPLE 1

2-Amino-6- $\{$ [4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxy-
carbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxymethyl $\}$
-4-pyrimidone

- 5 A solution of guanidine hydrochloride (0.20 g), ethyl 4- $\{$
[4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-
1,4-dihydropyridin-2-yl]methoxy $\}$ acetoacetate (0.99 g), and
1,5-diazabicyclo[4.3.0]non-5-ene (0.30 g) in ethanol (50 ml) was
stirred at room temperature for 5 days and then evaporated. The
10 residue was dissolved in chloroform and washed with 2M
hydrochloric acid, 10% aqueous sodium carbonate, and water, dried
(MgSO_4), and evaporated. The residual solid was recrystallised
from ethyl acetate to give the title compound (0.23 g), m.p.
185-187°C. Found: C, 56.08; H, 5.14; N, 11.33. $\text{C}_{23}\text{H}_{25}\text{ClN}_4\text{O}_6$
15 requires C, 56.50; H, 5.15; N, 11.46%.

EXAMPLE 2

2-Amino-6- $\{$ [4-(2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxy-
carbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxymethyl $\}$ -4-
pyrimidone

- 20 A solution of ethyl 4- $\{$ [4-(2,3-dichlorophenyl)-3-ethoxy-
carbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-
yl]methoxy $\}$ acetoacetate (0.74 g), 1,5-diazabicyclo[4.3.0]non-5-
ene (0.25 g) and guanidine hydrochloride (0.14 g) in ethanol (30
ml) was heated under reflux for 5.5 hours and then evaporated.
25 The residue was dissolved in chloroform and the solution washed
successively with water, 10% aqueous sodium carbonate solution and

water, dried over anhydrous magnesium sulphate and evaporated.

The residue was triturated with ethyl acetate and the resulting precipitate collected, washed thoroughly with diethyl ether, recrystallised from ethyl acetate/ethanol and dried to give the


5 title compound (0.20 g), m.p. 222-225°C. Found: C, 52.10;

H, 4.56; N, 10.54. $C_{23}H_{24}Cl_2N_4O_6$ requires C, 52.78; H, 4.62;

N, 10.70%.



EXAMPLES 3 - 12

The following compounds were prepared by the method of
10 process (c) as described in Example 1 or 2 by reacting the appropriate 4-(1,4-dihydropyridine-2-ylmethoxy)acetoacetate of formula (VI) with the appropriate amidine or guanidine of formula (VIa).



Example No.	R ¹	R ²	R ⁷	R ⁵	m.p. (°C)	Form Characterised	Analysis % (Theoretical in brackets) C H N
3	CH ₃	C ₂ H ₅	H	CH ₃	230-235 (decomp.)	hemihydrate	58.31 5.41 8.32 (58.00 5.48 8.45)
4	CH ₃	C ₂ H ₅	Cl	CH ₃	225-230	hydrate	53.64 4.70 7.91 (53.34 5.03 7.77)
5	C ₂ H ₅	CH ₃	Cl	NH ₂	220	hydrate	51.02 4.67 10.53 (51.02 4.47 10.35)
6	CH ₃	C ₂ H ₅	CF ₃	NH ₂	228-229	hemihydrate	50.96 4.41 10.02 (50.93 4.45 9.90)
7	C ₂ H ₅	CH ₃	CF ₃	NH ₂	220	hydrate	50.17 4.48 9.63 (50.13 4.56 9.74)
8	CH ₃	C ₂ H ₅	H	C ₆ H ₅	196-198	free base	62.71 5.18 7.50 (63.32 5.13 7.64)
9	CH ₃	C ₂ H ₅	Cl	N(CH ₃) ₂	219-222	free base	54.13 5.18 9.90 (54.45 5.12 10.16)

0132375

Example No.	R ¹	R ²	R ⁷	R ⁵	m.p. (°C)	Form Characterised	Analysis % (Theoretical in brackets) C H N
10	CH ₃	C ₂ H ₅	H		151-155	free base	57.02 5.56 10.08 (58.01 5.59 10.02)
11	CH ₃	C ₂ H ₅	Cl		147-150	free base	56.85 5.65 9.34 (56.85 5.45 9.47)

EXAMPLE 12

2-Amino-6- $\left\{ [4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl}] \text{methoxymethyl} \right\}$ -3-methyl-4-pyrimidone

- 5 A mixture of 2-amino-6- $\left\{ [4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl}] \text{methoxymethyl} \right\}$ -4-pyrimidone (0.52 g), iodomethane (0.14 g) and potassium carbonate (0.14 g) in dimethylformamide (20 ml) was stirred at room temperature for 4 days and then evaporated. The
- 10 residue was partitioned between chloroform and water and the organic layer washed twice with water, dried over anhydrous magnesium sulphate and evaporated. The residue was crystallised from ethyl acetate to give the title compound (0.23 g), m.p. 202-205°C. Found: C, 53.42; H, 4.87; N, 10.49. $\text{C}_{24}\text{H}_{26}\text{Cl}_2\text{N}_4\text{O}_6$
- 15 requires C, 53.64; H, 4.88; N, 10.43%.

EXAMPLE 13

- 6- $\left\{ [4-(2,3\text{-Dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl}] \text{methoxymethyl} \right\}$ -2,3-dimethyl-4-pyrimidone was prepared by the method described in
- 20 Example 12 using 6- $\left\{ [4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl}] \text{methoxymethyl} \right\}$ -2-methyl-4-pyrimidone as the starting material. The product had m.p. 153-158°C. Found: C, 55.66; H, 5.09; N, 7.97. $\text{C}_{25}\text{H}_{27}\text{Cl}_2\text{N}_3\text{O}_6$ requires C, 55.98; H, 5.07; N, 7.83%.


EXAMPLE 14


7- $\{$ 4-(2,3-Dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxymethyl $\}$ -5-hydroxy-imidazo[1,2-a]pyrimidine

- 5 A mixture of 2-aminoimidazolium sulphate (0.52 g), ethyl 4-
 $\{$ 4-(2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-
 methyl-1,4-dihydropyridin-2-yl]methoxy $\}$ acetoacetate (1.50 g) and
 1,5-diazabicyclo[4.3.0]non-5-ene (1.06 g) in ethanol (40 ml) was
 heated under reflux for 3.5 hours and then evaporated. The
10 residue was partitioned between chloroform and water and the
 organic layer washed three times with water, dried over anhydrous
 magnesium sulphate and evaporated. The residue was purified by
 chromatography on silica gel (3 g) using chloroform plus 0-1% by
 volume of methanol as eluant. Appropriate fractions were combined
15 and evaporated and the residue crystallised from ethyl acetate to
 give the title compound (0.28 g), m.p. 187-188°C. Found:
 C, 54.78; H, 4.64; N, 10.42. $C_{25}H_{24}Cl_2N_4O_6$ requires C, 54.85; H, 4.42;
 N, 10.24%.

EXAMPLE 15

- 20 The following compound was prepared by the method described
 in Example 14 using 2-iminopiperidine as the starting material.



Example No.	X	m.p. (°C)	Analysis % (Theoretical in Brackets)		
			C	H	N
15		177-8	54.13	5.18	9.90
			(54.45	5.12	10.16)

EXAMPLE 16

5 5-Σ[4-(2-Chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxymethylΣ-3-pyrazolone

A solution of ethyl 4-Σ[4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxyΣ acetoacetate (0.50 g) and hydrazine hydrate (0.50 g)

in ethanol (20 ml) was kept at room temperature for 4 days and then evaporated. The residue was purified by chromatography on silica gel (1.0 g) using dichloromethane plus 0-5% by volume of methanol as eluant. Appropriate fractions were combined and
5 evaporated and the residue crystallised from ethyl acetate to give the title compound (0.18 g), m.p. 207°C. Found: C, 57.11; H, 5.15; N, 9.19. $C_{22}H_{24}ClN_3O_6$ requires: C, 57.20; H, 5.24; N, 9.10%.

EXAMPLE 17

2-Amino-1-<2- $\{$ [4-(2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxy-
10 carbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy $\}$ ethyl>-4-imidazolone

A solution of 2- $\{$ 2-[(N-cyano-N-ethoxycarbonylmethyl)amino]-ethoxymethyl $\}$ -4-(2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine (0.70 g) in saturated
15 methanolic ammonia (10 ml) was stirred at room temperature for 16 hours and then evaporated. The residual solid was recrystallised from ethanol to give the title compound as a hemihydrate (0.46 g), m.p. 134-138°C. Found: C, 51.83; H, 5.27; N, 10.02.
 $C_{23}H_{26}Cl_2N_4O_6 \cdot 0.5 H_2O$ requires: C, 51.68; H, 5.06; N, 10.48%.

EXAMPLE 18

20 2-Amino-1-<2- $\{$ [4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy $\}$ -ethyl>-4-imidazolone was prepared by the method described in Example 17 using 4-(2-chlorophenyl)-2- $\{$ 2-[(N-cyano-N-ethoxy-
25 carbonylmethyl)amino]ethoxymethyl $\}$ -3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine as the starting material.

The product was characterised as its hemihydrate, m.p. 110-113°C.

Found: C,55.45; H,5.56; N,11.20. $C_{23}H_{27}ClN_4O_6 \cdot 0.5 H_2O$ requires:
C,55.20; H,5.60; N,11.20%.

EXAMPLE 19

5 1-<2- { [4-(2-Chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy } ethyl>imidazolidine-2,5-dione

A solution of 1-<2- { [4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy } ethyl>-
10 3-(ethoxycarbonylmethyl)urea (1.08 g) in methanol (1 ml) and .880 aqueous ammonia (2 ml) was stirred at room temperature for 16 hours and then evaporated. The residue was purified by chromatography on silica (2.5 g) using chloroform plus 0-4% v/v methanol as eluant. Appropriate fractions were combined and
15 evaporated and the residual solid recrystallised from a mixture of ethyl acetate and acetone to give the title compound (0.20 g), m.p. 140°C. Found: C,56.17; H,5.64; N,8.45. $C_{23}H_{26}ClN_3O_7$ requires C,56.15; H,5.33; N,8.54%.

EXAMPLE 20

20 1-<2- { [4-(2-Chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy } ethyl>imidazolidine-2,4-dione


Sodium hydride (60 mg; 80% dispersion in oil) was added to a suspension of 1-<2- { [4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy } ethyl>-
25

1-(methoxycarbonylmethyl)urea (0.52 g) in methanol (40 ml) and the mixture stirred at room temperature for one hour. The reaction mixture was evaporated and the residue partitioned between ethyl acetate and water. The organic layer was dried (MgSO_4) and evaporated and the residue triturated with diethyl ether. The resulting solid was collected, washed with diethyl ether and dried to give the title compound (0.41 g), m.p. 105-107°C (decomp.). Found: C, 56.20; H, 5.50; N, 8.36. $\text{C}_{23}\text{H}_{26}\text{ClN}_3\text{O}_7$ requires: C, 56.16; H, 5.33; N, 8.54%.





10




EXAMPLES 21-27

The following compounds were prepared by the method described in Example 20 using the appropriate starting materials of formula (VIII) and were characterised in the form indicated.



0132375

Example No.	R ⁷	X	Form characterised	m.p. (°C)	Analysis % (Theoretical in brackets) C H N
21	Cl		free base, hemihydrate	116-118	51.69 4.99 7.79 (51.60 4.86 7.85)
22	H		free base	166-167	56.65 5.59 8.25 (56.97 5.58 8.31)
23	Cl		free base, hemihydrate	115-116	52.44 4.94 7.51 (52.45 5.10 7.65)
24	H		free base, hydrate	132-134	54.73 5.51 11.17 (54.28 5.70 11.01)

Example No.	R ⁷	X	Form characterised	m.p. (°C)	Analysis % (Theoretical in brackets) C H N
25	Cl		hydrochloride, dihydrate	131-133	46.73 4.67 9.63 (46.54 5.23 9.44)
26	H		hydrochloride, dihydrate	138-139	49.64 5.44 10.21 (49.46 5.55 10.04)
27	H		free base, sesquihydrate	117-119	56.76 5.89 7.29 (56.66 6.10 7.32)

EXAMPLE 28

1-<2-Σ[4-(2-Chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxyΣethyl>-2-methyl-5-imidazolone

- 5 A solution of 2-(2-aminoethoxy)methyl-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine (1.02 g) and ethyl N-(ethoxycarbonylmethyl)acetimidate (0.43 g) in ethanol (20 ml) was stirred at room temperature for 16 hours and then evaporated. The residue was purified by chromatography on silica (10 g) using
10 dichloromethane plus 0-5% v/v methanol as eluant. Appropriate fractions were combined and evaporated to give the title compound (0.07 g), m.p. 167-169°C. Found: C,58.30; H,5.79; N,8.55. C₂₄H₂₈ClN₃O₆ requires: C,58.83; H,5.72; N,8.58%.

EXAMPLE 29


- 15 1-<2-Σ[4-(2-Chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxyΣethyl>-2-pyridone

- A solution of 1-(2-hydroxyethyl)-2-pyridone (6.15 g) in tetrahydrofuran (30 ml) was added over 45 minutes to a suspension of sodium hydride (3.0 g; 80% dispersion in oil) in
20 tetrahydrofuran (30 ml). The mixture was stirred at room temperature for one hour, treated with a solution of ethyl 4-chloroacetoacetate (8.23 g) in tetrahydrofuran (20 ml) dropwise over 3 hours and then stirred at room temperature for 16 hours. The reaction mixture was treated with ethanol (10 ml), poured into
25 concentrated hydrochloric acid (10 ml) and extracted into ethyl acetate. The ethyl acetate extract was dried (MgSO₄) and



evaporated to give ethyl 4-[2-(2-pyridon-1-yl)ethoxy]acetoacetate as a brown oil which was shown by n.m.r. to be 75% pure. A solution of this crude product (5.4 g) and ammonium acetate (1.50 g) in ethanol (20 ml) was heated at 50°C for 30 minutes, treated with methyl 2-(2-chlorobenzylidene)acetoacetate (3.57 g) and the mixture heated under reflux for 3 hours. The reaction mixture was evaporated and the residue dissolved in toluene, washed with 2M hydrochloric acid, dried (MgSO_4) and evaporated. The residue was purified by chromatography on silica (50 g) using petroleum ether (b.p. 40-60°C) plus 70-100% v/v dichloromethane as eluant. Appropriate fractions were combined and evaporated to give the title compound (0.15 g), m.p. 114-115°C. Found: C, 61.66; H, 5.55; N, 5.75. $\text{C}_{25}\text{H}_{27}\text{ClN}_2\text{O}_6$ requires C, 61.37; H, 5.65; N, 5.70%.

EXAMPLES 30 AND 31

The following compounds were prepared by the method described in Example 29 using the appropriate starting materials of formulae (IV) and (V) and were characterised in the form indicated.



0132375

Example No.	X	Form characterised	m.p. (°C)	Analysis % (Theoretical in brackets) C H N
30		dihydrochloride hydrate	90	49.95 5.93 7.31 (49.45 5.88 7.21)
31		free base	152-153	53.06 4.86 11.42 (53.47 4.82 11.55)

EXAMPLE 32

1-(2- Σ [4-(2-Chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy Σ ethyl)-4-ethoxycarbonyl-5-hydroxy-(1H)-1,2,3-triazole

5 A mixture of 2-(2-azidoethoxy)methyl-4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine (2.17 g), diethyl malonate (4.00 g) and sodium hydride (0.75 g; 80% dispersion in oil) in ethanol (80 ml) was heated under reflux for 16 hours and then evaporated. The residue was partitioned
10 between 0.1 M hydrochloric acid and ethyl acetate and the organic layer was washed with water, dried (Na_2SO_4) and evaporated. The residue was triturated with ether and the resulting solid collected and purified by chromatography on silica (30 g) using dichloromethane plus 0-5% v/v methanol as eluant. Appropriate
15 fractions were combined and evaporated to give the title compound (0.89 g), m.p. 121-124°C. Found: C, 55.01; H, 5.49; N, 10.07. $\text{C}_{25}\text{H}_{29}\text{ClN}_4\text{O}_8$ requires: C, 54.69; H, 5.32; N, 10.21%.

EXAMPLE 33

2-(2- Σ [4-(2,3-Dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy Σ ethyl)-1-oxo-1,3-dihydropyrrolo[2,3-b]pyridine

20 A mixture of 2-(2-aminoethoxy)methyl-4-(2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine (0.50 g), 2-bromomethyl-3-ethoxycarbonylpyridine (0.31 g) and
25 potassium carbonate (0.56 g) in acetonitrile (30 ml) was heated under reflux for 41 hours, filtered and evaporated. The residue

was purified by chromatography on silica (10 g) using dichloromethane plus 0-5% v/v methanol as eluant. Appropriate fractions were combined and evaporated and the residue triturated with ether. The resulting solid was collected, washed with ether
5 and dried to give the title compound (0.22 g), m.p. 147-149°C.
Found: C, 57.96; H, 5.07; N, 7.47. $C_{27}H_{27}Cl_2N_3O_6$ requires C, 57.86; H, 4.86; N, 7.50%.

EXAMPLE 34

5-Cyano-1-(2-[4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxy-
10 carbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy}ethyl)-3-
methylpyrimidine-2,4-dione

A solution of 2-(2-aminoethoxy)methyl-4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine (0.82 g) and 2-cyano-N-ethoxycarbonyl-2-ethoxymethylene-N-
15 methylacetamide (0.5 g) in methanol (2 ml) containing one drop of triethylamine was heated under reflux for 2 hours, diluted with toluene (5 ml) and evaporated. The residue was purified by chromatography on silica (5 g) using toluene plus 0-100% v/v chloroform as eluant. Appropriate fractions were combined and
20 evaporated and the residue crystallised from a mixture of ethyl acetate and diethyl ether. The resulting solid was collected and recrystallised from a mixture of acetone and diisopropyl ether to give the title compound (0.25 g), m.p. 155-156°C. Found:
C, 57.46; H, 5.09; N, 10.13. $C_{26}H_{27}ClN_4O_7$ requires: C, 57.50;
25 H, 5.03; N, 10.32%.

EXAMPLE 35

1-<2-Σ[4-(2-Chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxyΣethyl>maleimide

A solution of 2-(2-aminoethoxy)methyl-4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine (0.82 g) and maleic anhydride (0.25 g) in acetonitrile (2 ml) was heated under reflux for 50 minutes, treated with acetic anhydride (1 ml), heated under reflux for a further one hour, diluted with methanol (2 ml), heated under reflux for a further 20 minutes and evaporated. The residue was partitioned between ethyl acetate and 10% aqueous sodium carbonate and the organic layer dried (Na₂CO₃) and evaporated. The residue was purified by chromatography on silica (3 g) using toluene plus 0-100% v/v chloroform as eluant. Appropriate fractions were combined and evaporated and the residue triturated with diethyl ether. The resulting solid was collected and recrystallised from a mixture of acetone and diisopropyl ether to give the title compound (0.18 g), m.p. 140-141°C. Found: C, 58.77; H, 5.16; N, 5.85. C₂₄H₂₅ClN₂O₇ requires: C, 58.95; H, 5.15; N, 5.73%.

EXAMPLE 36

Tablets are compounded from the following ingredients:

	<u>mg/tablet</u>
Product of any one of Examples	10
Dicalcium phosphate	120
25 Magnesium stearate	1.8
Sodium lauryl sulphate	0.2

The ingredients are thoroughly blended, compressed, granulated and recompressed to tablets of the desired size.

EXAMPLE 37

Capsules are compounded from the following ingredients:

	<u>mg/capsule</u>
5	
Product of any one of Examples	10
Maize starch	127
Cellulose (microcrystalline)	127
Magnesium stearate	5.4
10	
Sodium lauryl sulphate	0.6

The ingredients are thoroughly blended, then filled into hard gelatine capsules of the appropriate size to contain the ingredients.

Preparation 1

15 Ethyl 4- Σ [4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy $\}$ acetoacetate

A solution of 2- Σ [4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy $\}$ acetic acid (4.24 g) in tetrahydrofuran (60 ml) was treated with
 20 ice-cooling with carbonyldiimidazole (1.70 g) and the mixture stirred at 0°C for 2.25 hours. This solution was added dropwise over 10 minutes to a solution of the sodium salt of 2,2-dimethyl-1,3-dioxane-4,6-dione in dimethylformamide (prepared by stirring a mixture of 2,2-dimethyl-1,3-dioxane-4,6-dione (1.44 g) and sodium

hydride (0.30 g; 80% dispersion in oil) in dimethylformamide (30 ml) with ice-cooling for 30 minutes) and the mixture stirred for 16 hours at room temperature and then evaporated. The residue was dissolved in dichloromethane, washed twice with 1M hydrochloric acid and water, dried (MgSO_4), and evaporated. A solution of the residual oil (6 g) in ethanol (50 ml) was heated under reflux for 6 hours and evaporated. The residue was dissolved in diethyl ether, washed twice with 10% aqueous sodium carbonate and once with saturated aqueous sodium chloride, dried (MgSO_4), and evaporated. The residue was purified by chromatography on silica (30 g) using hexane plus 5-15% v/v ethyl acetate as eluant. Appropriate fractions were combined and evaporated and the residue crystallised from ethanol to give the title compound (1.5 g), m.p. 88-89°C. Found: C, 58.18; H, 5.73; N, 2.85. $\text{C}_{24}\text{H}_{28}\text{ClNO}_8$ requires: C, 58.36; H, 5.71; N, 2.84%.

Preparation 2


Ethyl 4- $\left\{ \left[4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl} \right] \text{methoxy} \right\}$ acetoacetate

A solution of carbonyl diimidazole (5.20 g) and 2- $\left\{ \left[4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl} \right] \text{methoxy} \right\}$ acetic acid (14.00 g) in dichloromethane (200 ml) was stirred at room temperature under nitrogen for 2 hours and then added to a solution of pyridine (2.40 g) and 2,2-dimethyl-1,3-dioxane-4,6-dione (4.56 g) in dichloromethane (200 ml) over 8 minutes. The mixture was stirred at room temperature for 2.5 hours, washed successively with water,

ice-cold 2.5 M hydrochloric acid and saturated brine, dried over anhydrous magnesium sulphate and evaporated. The residue was dissolved in ethanol (200 ml) and the solution heated under reflux for 2.75 hours and evaporated. The residue was triturated with
 5 diethyl ether and the resulting solid collected washed with diethyl ether and dried to give the title compound (9.0 g), m.p. 99-102°C.

Preparations 3-6

The following compounds were prepared by the method described
 10 in Preparations 1 and 2 from the appropriate 2-(dihydropyridin-2-ylmethoxy)acetic acid.



Preparation No.	R ¹	R ²	R ⁷	m.p. (°C)	Analysis % (Theoretical in brackets) C H N		
3	CH ₃	CH ₂ CH ₃	Cl	99-102	characterised by 'H-n.m.r.		
4	CH ₂ CH ₃	CH ₃	Cl	134-7	54.44 (54.55)	5.21 5.15	2.64 2.65
5	CH ₃	CH ₂ CH ₃	CF ₃	96	characterised by 'H-n.m.r.		
6	CH ₂ CH ₃	CH ₃	CF ₃	125-8	53.04 (53.43)	4.80 4.84	2.44 2.49

Preparation 7

2- $\{$ 2-[(N-Cyano-N-ethoxycarbonylmethyl)amino]ethoxymethyl $\}$ -4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine

10 Cyanogen bromide (0.35 g) was added to a stirred solution of
4-(2-chlorophenyl)-3-ethoxycarbonyl-2- $\{$ 2-[(N-ethoxycarbonyl-
methyl)amino]ethoxymethyl $\}$ -5-methoxycarbonyl-6-methyl-1,4-
dihydropyridine (1.25 g) in chloroform (10 ml) containing sodium
hydrogen carbonate (0.3 g). The mixture was stirred at room
15 temperature for 20 hours and then evaporated. The residue was
trituated with diethyl ether and the resulting solid collected,
washed with diethyl ether and dried to give the title compound as
a hemihydrate (0.81 g), m.p. 132-134°C. Found: C, 56.96; H, 5.66;
N, 7.94. C₂₅H₃₀ClN₃O₇ · 0.5 H₂O requires: C, 56.76; H, 5.86; N, 7.94%.

Preparation 8

2- Σ 2-[(N-Cyano-N-ethoxycarbonylmethyl)amino]ethoxymethyl
 -4-(2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-
 methyl-1,4-dihydropyridine was prepared by the method described in
 5 Preparation 7 using 4-(2,3-dichlorophenyl)-3-ethoxycarbonyl-2-
 2-[(N-ethoxycarbonylmethyl)amino]ethoxymethyl Σ -5-methoxycarbonyl-
 6-methyl-1,4-dihydropyridine as the starting material. The
 product was obtained as a hemihydrate, m.p. 148-150°C. Found:
 C, 53.08; H, 5.04; N, 7.46. $C_{25}H_{29}Cl_2N_3O_7 \cdot 0.5 H_2O$ requires:
 10 C, 53.28; H, 5.33; N, 7.46%.

Preparation 9

1-<2- Σ [4-(2-Chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-
 methyl-1,4-dihydropyridin-2-yl]methoxy Σ ethyl>-3-(cyanomethyl)urea

A solution of 4-(2-chlorophenyl)-3-ethoxycarbonyl-2- Σ 2-[N-
 15 (1-imidazolylcarbonyl)amino]ethoxymethyl Σ -5-methoxycarbonyl-6-
 methyl-1,4-dihydropyridine (0.98 g), aminoacetonitrile
 hydrochloride (0.30 g) and N-methylmorpholine (0.44 g) was
 stirred at room temperature for 20 hours and then evaporated. The
 residue was purified by chromatography on silica (10 g) using
 20 dichloromethane plus 0-5% v/v methanol as eluant. Appropriate
 fractions were combined and evaporated and the residual solid
 recrystallised from ethanol to give the title compound as a
 hemihydrate (0.50 g), m.p. 145-147°C. Found: C, 55.34; H, 5.84;
 N, 11.33. $C_{23}H_{27}ClN_4O_6 \cdot 0.5 H_2O$ requires: C, 55.25; H, 5.60;
 25 N, 11.21%.

Preparation 10

4-(2-Chlorophenyl)-3-ethoxycarbonyl-2-~~2~~-[N-(1-imidazolyl-
carbonyl)amino]ethoxymethyl-5-methoxycarbonyl-6-methyl-1,4-
dihydropyridine

5 A solution of 2-(2-aminoethoxy)methyl-4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine (20.4 g), carbonyldiimidazole (8.9 g) and N-methylmorpholine (20 ml) in tetrahydrofuran (500 ml) was stirred at room temperature for 2 hours and then evaporated. The residue was partitioned
 10 between ethyl acetate and water and the organic layer washed with water, dried (Na_2SO_4), and evaporated. The residual solid was washed with diethyl ether and dried to give the title compound (16.6 g), m.p. 149-151°C. Found: C, 57.35; H, 5.54; N, 11.22. $\text{C}_{24}\text{H}_{27}\text{ClN}_4\text{O}_6$ requires: C, 57.54; H, 5.43; N, 11.18%.

Preparation 11

15 1-~~2~~-[4-(2-Chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-
methyl-1,4-dihydropyridin-2-yl]methoxy ethyl-1-(cyanomethyl)urea

 A mixture of 2-(2-aminoethoxy)methyl-4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine
 20 (4.08 g), chloroacetonitrile (1.20 g), and potassium carbonate (1.20 g) in acetonitrile (40 ml) was heated under reflux for 16 hours and then evaporated. The residue was triturated with diethyl ether and the resulting solid dried and purified by chromatography on silica (40 g) using dichloromethane plus 0-5%
 25 v/v methanol as eluant. Appropriate fractions were combined and evaporated to give crude 4-(2-chlorophenyl)-2-[2-(N-


cyanomethyl)amino-ethoxy]methyl-3-ethoxycarbonyl-5-methoxy-
 carbonyl-6-methyl-1,4- dihydropyridine (2.0 g). Acetic acid (0.5
 g) was added to a solution of the above product (1.0 g) and
 potassium cyanate (0.35 g) in water (15 ml) and dioxane (15 ml)
 5 and the mixture stirred at room temperature for 16 hours and then
 evaporated. The residue was partitioned between ethyl acetate and
 water and the organic layer dried (Na_2SO_4) and evaporated. The
 residue was purified by chromatography on silica (10 g) using
 dichloromethane plus 0-5% v/v methanol as eluant. Appropriate
 10 fractions were combined and evaporated to give the title compound
 (53 mg), m.p. 190-191°C. Found: C, 56.34; H, 5.80; N, 10.99.
 $\text{C}_{23}\text{H}_{27}\text{ClN}_4\text{O}_6$ requires: C, 56.27; H, 5.50; N, 11.42%.

Preparation 12

1-<2- $\left\{ \begin{array}{l} \text{[4-(2,3-Dichlorophenyl)-3-ethoxycarbonyl-5-methoxy-} \\ \text{15 carbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy} \end{array} \right\} \text{ethyl}>$ -1-
 (cyanomethyl)urea was prepared by the method described in
 Preparation 11 using 2-(2-aminoethoxy)methyl-4-(2,3-dichloro-
 phenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-
 dihydropyridine as starting material. The product was obtained as
 20 a foam. Rf (silica; chloroform, methanol, ammonium hydroxide;
 80:20:1) 0.7.

CLAIMS

1. A 1,4-dihydropyridine compound having the formula:-



and pharmaceutically acceptable salts thereof;

wherein R is aryl or heteroaryl;

R^1 and R^2 are each independently $\text{C}_1\text{-C}_4$ alkyl or 2-methoxyethyl;

X is a 5 or 6 membered nitrogen-containing heterocyclic ring which is substituted with one or more hydroxyl or oxo groups and which may optionally be fused to a further 5 or 6 membered nitrogen-containing heterocyclic ring, and which may optionally be additionally substituted in the heterocyclic ring or further fused heterocyclic ring by one or more $\text{C}_1\text{-C}_4$ alkyl, phenyl, CN, $\text{N}(\text{R}^3)_2$, $(\text{CH}_2)_m\text{CO}_2\text{R}^3$ or $(\text{CH}_2)_m\text{CON}(\text{R}^3)_2$ groups, wherein each R^3 is independently H or $\text{C}_1\text{-C}_4$ alkyl or the two groups R^3 are taken together with the nitrogen atom to which they are attached to form a pyrrolidinyl, piperidino, morpholino, piperazinyl or N-($\text{C}_1\text{-C}_4$ alkyl)-piperazinyl group and m is 0 or 1;

Y is $-(\text{CH}_2)_n-$, $-\text{CH}_2\text{CH}(\text{CH}_3)-$ or $-\text{CH}_2\text{C}(\text{CH}_3)_2-$;

and n is 1 to 3 when X is linked to Y by a ring carbon atom,
or 2 or 3 when X is linked to Y by a ring nitrogen atom.

2. A compound according to claim 1, wherein R is "aryl" and "aryl" is phenyl or phenyl substituted by one or two substituents each independently selected from nitro, halo, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, trifluoromethyl, and cyano, or is 1- or 2-naphthyl; or wherein R is "heteroaryl" and "heteroaryl" is benzofuranyl; benzothienyl; pyridyl optionally substituted by methyl, methylthio, cyano or halo; quinolyl; benzoxazolyl; benzothiazolyl; furyl; pyrimidinyl; thiazolyl; 2,1,3-benzoxadiazol-4-yl; 2,1,3-benzothiadiazol-4-yl or thienyl optionally monosubstituted by halo or C₁-C₄ alkyl.

3. A compound according to claim 2 wherein R is 2-chlorophenyl or 2,3-dichlorophenyl.

4. A compound according to any one of claims 1 to 3 wherein X is 2-, 4- or 5-imidazolone, 3-pyrazolone, pyrrolone-2,5-dione, imidazolidine-2,4-dione, 2-pyridone, 3-piperazinone, 4-pyrimidone, pyrimidine-2,4-dione, 5-hydroxy-(1H)-1,2,3-triazole or 1-oxo-1,3-dihydropyrrolo- [2,3b]pyridine, any of which may optionally be substituted by amino, lower alkyl, carbamoyl or CN.

5. A compound according to claim 4 wherein X is 1-(2-amino-4-imidazolone), 1-(imidazolidine-2,4-dione), 1-(4-amino-2-imidazolone), 2-(1-oxo-1,3-dihydropyrrolo[2,3b]-pyridine), 6-(2-amino-4-pyrimidone) or 6-(2,3-dimethyl-4-pyrimidone).

6. A compound according to any one of claims 1 to 5 wherein R^1 is CH_3 and R^2 is C_2H_5 .

7. A compound according to any one of claims 1 to 6 wherein Y is $-(CH_2)_n-$ and n is 1 or 2.

8. A compound according to claim 1 being

2-Amino-1- $\left\{2-[4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl}]methoxy\right\}$ ethyl-4-imidazolone;

1- $\left\{2-[4-(2,3\text{-Dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl}]methoxy\right\}$ ethyl-imidazolidine-2,4-dione;

4-Amino-1- $\left\{2-[4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl}]methoxy\right\}$ ethyl-2-imidazolone;

2- $\left\{2-[4-(2,3\text{-Dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl}]methoxy\right\}$ ethyl-1-oxo-1,3-dihydropyrrolo[2,3-b]pyridine;

6- $\left\{[4-(2,3\text{-Dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl}]methoxymethyl\right\}$ -2,3-dimethyl-4-pyrimidone, or


2-Amino-6- $\left\{ \begin{array}{l} [4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-} \\ \text{methoxy-carbonyl-6-methyl-1,4-dihydropyridin-2-yl}] \text{methoxy-} \\ \text{methyl} \end{array} \right\}$ -4-pyrimidone

9. A pharmaceutical composition comprising a compound of the formula (I) as claimed in any one of the preceding claims, or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable diluent or carrier.

10. A compound of the formula (I) as claimed in any one of claims 1 to 8, or a pharmaceutically acceptable salt thereof, for use in the treatment of cardiovascular conditions, including use in the prevention or treatment of cardiac conditions, or use as an antihypertensive, in man.

CLAIMS FOR THE CONTRACTING STATE: AT

1. A Process for preparing 1,4-dihydropyridine compounds having the formula:-



and pharmaceutically acceptable salts thereof;

wherein R is aryl or heteroaryl;


R¹ and R² are each independently C₁-C₄ alkyl or 2-methoxyethyl;

X is a 5 or 6 membered nitrogen-containing heterocyclic ring which is substituted with one or more hydroxyl or oxo groups and which may optionally be fused to a further 5 or 6 membered nitrogen-containing heterocyclic ring, and which may optionally be additionally substituted in the heterocyclic ring or further fused heterocyclic ring by one or more C₁-C₄ alkyl, phenyl, CN, N(R³)₂, (CH₂)_mCO₂R³ or (CH₂)_mCON(R³)₂ groups, wherein each R³ is independently H or C₁-C₄ alkyl or the two groups R³ are taken together with the nitrogen atom to which they are attached to form a pyrrolidinyl, piperidino, morpholino, piperazinyl or N-(C₁-C₄ alkyl)-piperazinyl group and m is 0 or 1;

Y is -(CH₂)_n-, -CH₂CH(CH₃)- or -CH₂C(CH₃)₂-;


and n is 1 to 3 when X is linked to Y by a ring carbon atom,
or 2 or 3 when X is linked to Y by a ring nitrogen atom.
which process comprises:

(a) reacting the following compounds:




wherein R, R^1 , R^2 , X and Y are as previously defined, or

(b) reacting together compounds of the formula:



wherein R, R^1 , R^2 , X and Y are as previously defined, or

(c) for preparing the compounds of formula (I) wherein X is a 4-pyrimidone or 3-pyrazolone group, reacting a dihydropyridine ketoester of formula:




wherein R, R¹, R² and Y are as previously defined and R⁴ is C₁-C₄ lower alkyl; with a compound of the formula:




wherein R⁵ is N(R³)₂, C₁-C₄ alkyl or phenyl, where R³ is as previously defined, to give compounds wherein X is a 2-substituted-4-pyrimidone group; or with hydrazine to give compounds wherein X is a 3-pyrazolone group, and optionally alkylating with a C₁-C₄ alkyl halide to introduce one or further C₁-C₄ alkyl groups into X; or

(d) for preparing the compounds of formula (I) wherein X is a 1-(2-amino-4-imidazolone) group reacting a compound of the formula:



wherein R, R¹, R², R⁴ and Y are as previously defined, with ammonia; or


(e) for preparing the compounds of formula (I) wherein X is a 1-(imidazolidine-2,4-dione) group or a 1-(4-amino-2-imidazolone) group which may be substituted on the 3- or 5-positions with lower alkyl groups, reacting a compound of the formula:



0132375


wherein R, R¹, R² and Y are as previously defined, each R⁶ is independently H or C₁-C₄ alkyl and Q is CO₂(C₁-C₄ alkyl) or CN; with a base; or

(f) for preparing the compounds of formula (I) wherein X is a 1-(5-amino-2-imidazolone) or a 1-(imidazolidine-2,5-dione) group, reacting a compound of the formula:




wherein R, R¹, R², Q and Y are as previously defined, with ammonia or with a base as appropriate; or

(g) for preparing the compounds of formula (I) wherein X is a 2-methyl-5-imidazolone group, reacting the corresponding 2-(aminoalkoxymethyl)-1,4-dihydropyridine with ethyl N-(ethoxycarbonylmethyl)acetimidate according to the following reaction scheme:



wherein R, R¹, R² and Y are as previously defined; or

(h) for preparing the compounds of formula (I) wherein X is a 4-ethoxycarbonyl- 5-hydroxy-(1H)-1,2,3-triazole group, reacting the corresponding 2-azidoalkoxymethyl-1,4-dihydropyridine of formula:



wherein R, R¹, R² and Y are as previously defined with diethyl malonate in the presence of a base; or

i) for preparing the compounds of formula (I) wherein X is a 2-(1-oxo-1,3- dihydropyrrolo[2,3-b]pyridine) group, reacting the 2-(aminoalkoxymethyl)-1,4-dihydropyridine of formula (X) wherein R, R¹, R² and Y are as previously defined, with 2-bromomethyl-3-ethoxycarbonylpyridine; or

j) for preparing the compounds of formula (I) wherein X is a 1-(5-cyano-3- methylpyrimidine-2,4-dione) group, reacting the 2-(aminoalkoxymethyl)-1,4-dihydropyridine of formula (X) wherein R, R¹, R² and Y are as previously defined, with 2-cyano-N-ethoxycarbonyl-2-ethoxymethylene-N- methylacetamide in a lower alkanol solvent in the presence of a trace of base; or

k) for preparing the compounds of formula (I) wherein X is a 1-(dihydro-pyrrol-2,5-dione) group, reacting the 2-(aminoalkoxymethyl)-1,4-dihydropyridine of formula (X) with maleic anhydride in an inert organic solvent; and in each case optionally forming a pharmaceutically acceptable salt of the product.

2. A process according to claim 1, wherein R is "aryl" and "aryl" is phenyl or phenyl substituted by one or two substituents each independently selected from nitro, halo, C_1-C_4 alkyl, C_1-C_4 alkoxy, hydroxy, trifluoromethyl, and cyano, or is 1- or 2-naphthyl; or wherein R is "heteroaryl" and "heteroaryl" is benzofuranyl; benzothienyl; pyridyl optionally substituted by methyl, methylthio, cyano or halo; quinolyl; benzoxazolyl; benzothiazolyl; furyl; pyrimidinyl; thiazolyl; 2,1,3-benzoxadiazol-4-yl; 2,1,3-benzothiadiazol-4-yl or thienyl optionally monosubstituted by halo or C_1-C_4 alkyl.

3. A process according to claim 2 wherein R is 2-chlorophenyl or 2,3-dichlorophenyl.

4. A process according to any one of claims 1 to 3 wherein X is 2-, 4- or 5-imidazolone, 3-pyrazolone, pyrrolone-2,5-dione, imidazolidine-2,4-dione, 2-pyridone, 3-piperazinone, 4-pyrimidone, pyrimidine-2,4-dione, 5-hydroxy-(1H)-1,2,3-triazole or 1-oxo-1,3-dihydropyrrolo- [2,3b]pyridine, any of which may optionally be substituted by amino, lower alkyl, carbamoyl or CN.

5. A process according to claim 4 wherein X is 1-(2-amino-4-imidazolone), 1-(imidazolidine-2,4-dione), 1-(4-amino-2-imidazolone), 2-(1-oxo-1,3-dihydropyrrolo[2,3b]-pyridine), 6-(2-amino-4-pyrimidone) or 6-(2,3-dimethyl-4-pyrimidone).

6. A process according to any one of claims 1 to 5 wherein R^1 is CH_3 and R^2 is C_2H_5 .

7. A process according to any one of claims 1 to 6 wherein Y is $-(CH_2)_n-$ and n is 1 or 2.

8. A process according to claim 1 wherein the compound of formula (I) is:

2-Amino-1- $\{2-[4-(2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy\}$ ethyl-4-imidazolone;

1- $\{2-[4-(2,3-Dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy\}$ ethyl-imidazolidine-2,4-dione;

4-Amino-1- $\{2-[4-(2,3-dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy\}$ ethyl-2-imidazolone;

2- $\{2-[4-(2,3-Dichlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy\}$ ethyl-1-oxo-1,3-dihydropyrrolo[2,3-b]pyridine.

6- $\left\{ [4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxy-} \right.$
carbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxymethyl $\left. \right\}$
-2,3-dimethyl-4-pyrimidone, or

2-Amino-6- $\left\{ [4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-} \right.$
methoxy-carbonyl-6-methyl-1,4-dihydropyridin-2-yl]methoxy-
methyl $\left. \right\}$ -4-pyrimidone

9. A process as claimed in claim 1 part (c) which comprises reacting a C_1-C_4 alkyl ester of 4- $\left\{ [4-(2,3\text{-dichlorophenyl})-3\text{-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridin-} \right.$
2-yl]methoxy $\left. \right\}$ acetoacetate and guanidine or a salt thereof in an organic solvent in the presence of a base.

10. A process as claimed in claim 9 wherein the solvent is ethanol and the reaction is heated at reflux temperature.